

Moduli Spaces of Instantons on Noncommutative 4-Manifolds II

Noriaki HAYAKAWA and Hiroshi TAKAI

Department of Mathematics,
Tokyo Metropolitan University

Abstract

Studied are moduli spaces of self dual connections on noncommutative 4-manifolds, especially deformation quantization of compact toric Riemannian 4-manifolds. Then such moduli spaces of irreducible modules associated with highest weights of compact connected semisimple Lie groups are smooth manifolds with dimension determined by their weights. This may be viewed as a generalization of Atiyah-Hitchin-Singer's classical result, Landi-Suijlekom's noncommutative 4-sphere case and the recent result of the second authors.

§1. Introduction Among many important topics of Yang-Mills theory, Atiyah-Hitchin-Singer [AHS] showed that the moduli spaces of irreducible self-dual connections of principal bundles over compact spin 4-manifolds are smooth manifolds with their dimension determined in terms of topological invariants related to ambient bundles. Recent development of noncommutative geometry serves many powerful devices toward determining a noncommutative analog of their result cited above. For example, Connes-Landi[CL] showed an existence of an isospectral deformation of compact spin Riemannian manifolds whose isometry groups have at least 2-torus subgroup. Related with this results, Connes-Devois-Violette [CV] observed that a deformation quantization of manifolds could be viewed as the fixed point algebra of the set of all smooth functions from manifolds to noncommutative tori under certain action of tori. Succeedingly, Landi-Suijlekom [LS] computed concerning the noncommutative 4-sphere, the index of the Dirac operator on certain spin modules as well as the dimension of the instanton moduli space of an irreducible module. Recently, the second author [T] verified that given noncommutative principal bundles based on compact spin toric Riemannian 4-manifolds and compact connected semisimple Lie groups and their associated noncommutative smooth sections with highest weights, their instanton moduli spaces are locally smooth manifolds with their dimension determined by their weights and the ambient bundles and as a corollary if the spaces of noncommutative anti-self-dual harmonic 2-forms vanish for all their self-dual connections, they are smooth manifolds with more explicit dimensions.

In this paper, we show that given a principal bundle based on compact toric Riemannian 4-manifold and a compact connected Lie group and its associated noncommutative smooth sections with a highest weight, its instanton moduli space is a smooth manifolds with its dimension determined by the highest weight and the ambient bundles. This result is used to compute the concrete example associated with the Hopf bundle over the complex projective 2-space.

§2. Noncommutative Yang-Mills Theory

In this section, we review briefly both commutative Yang-Mills theory due to Atiyah-Hitchin-Singer [AHS] and the noncommutative Yang-Mills theory due to Landi-Suijlekom [LS]. Let M be a compact oriented toric Riemannian 4-manifold, G a compact connected Lie group, and let P a principal G -bundle over M . Suppose there exist a smooth action φ from 2-torus T^2 into the isometry group $Iso(M, g)$ of M with respect to its Riemannian metric g . By Connes-

Devois-Violette [CV], let M_θ be a deformation quantization of M along θ as Frechet $*$ -algebra. Actually, there exists a quantization map L_θ from $C^\infty(M)$ onto M_θ such that $L_\theta(f \cdot g) = L_\theta(f) \times_\theta L_\theta(g)$. Then it is identified with the fixed point algebra of the set of all smooth maps from M to the noncommutative 2-torus T_θ^2 under the diagonal tensor action $\varphi \otimes \alpha^{-1}$ of T^2 where α is the gauge action of T^2 on T_θ^2 . Since M is 4-dimensional, its Hodge $*$ -operation on the Grassmann algebra $\Omega(M)$ of all forms of M could be shifted on the all forms $\Omega(M_\theta)$ of M_θ , which is denoted by $*_\theta$. Let us take a noncommutative principal bundle as follows: let

$$G \xrightarrow{\varrho} N \xrightarrow{\pi} M$$

be a principal G -bundle over M . Suppose there exists a smooth action $\tilde{\varphi}$ of a covering group $\tilde{T}^2$ of T^2 to $\text{Iso}(N, \pi^*(g))$ commuting with ϱ , then it follows from [LS] that there exists a smooth action $\tilde{\varrho}$ of G on $N_{\tilde{\theta}}$ such that

$$G \xrightarrow{\tilde{\varrho}} N_{\tilde{\theta}} \xrightarrow{\pi_\theta} M_\theta$$

is a noncommutative G -bundle over M_θ where $\pi^*(g)$ means the pull back Riemannian metric of g on N under π . Let σ be a highestweight of G with respect to its maximal torus T and V_σ its irreducible G -module. We identify σ with its associated irreducible representation of G . Let Ξ_σ be the fixed point algebra of $N_{\tilde{\theta}} \otimes V_\sigma$ under the diagonal tensor action $\tilde{\varrho} \otimes \sigma^{-1}$ of G . Then it is a finitely generated projective irreducible right M_θ -module, and there exists a natural number n and a projection P_σ in $M_n(M_\theta)$ such that $\Xi_\sigma = P_\sigma(M_\theta^n)$. We now take the Grassman connection $\nabla_\sigma = P_\sigma d_\theta^n$ of Ξ_σ where d_θ is the canonical outer derivative of $\Omega(M_\theta)$ derived by the ordinary outer derivative d of $\Omega(M)$. We have nothing to know at the moment whether ∇_σ is self-dual or anti selfdual although this is the case in the special set up (for instance [CDV],[LS]). Let us denote by $\mathcal{C}_+(\Xi_\sigma)$ the set of all self-dual compatible connections of Ξ_σ . In what follows, we treat the case that $\mathcal{C}_+(\Xi_\sigma)$ is non empty although we have nothing to check their existence for a given principal G -bundle over M and an irreducible representation of G . According to [LS], the set $\mathcal{C}_{\text{YM}}(\Xi_\sigma)$ of all Yang-Mills connections of Ξ_σ contains the sum of $\mathcal{C}_+(\Xi_\sigma)$

§3. Geometry of Noncommutative Instantons

In this section, we analyze a geometric structure of $\mathcal{C}_+(\Xi_\sigma)$ defined in the previous section under the case where they are non empty. Let $\nabla_\sigma \in \mathcal{C}_+(\Xi_\sigma)$ for an irreducible representation σ of G on a

finite dimensional $\mathbb{R}$ -vector space V_σ . Let P_- be the projection from $\Omega^2(M_\theta, \hat{\Xi}_\sigma)$ onto $\Omega_-^2(M_\theta, \hat{\Xi}_\sigma)$, where $\hat{\Xi}_\sigma = \text{End}_{M_\theta}(\Xi_\sigma)$. Since $\text{End}_{M_\theta}(\Xi_\sigma)$ is isomorphic to $(N_{\hat{\theta}} \otimes \mathcal{L}(V_\sigma))^{\hat{\theta} \otimes \text{Ad}(\sigma)(G)}$, then its skew-adjoint part $\hat{\Xi}_\sigma^{sk}$ is identified with $(N_{\hat{\theta}}^s \otimes \mathcal{L}^{sk}(V_\sigma))^{\hat{\theta} \otimes \text{Ad}(\sigma)(G)}$, where $\mathcal{L}(V_\sigma)$ is the set of all $\mathbb{C}$ -linear maps on V_σ , $\mathcal{L}^{sk}(V_\sigma)$ its skew-adjoint part, and $N_{\hat{\theta}}^s$ is the set of all self-adjoint elements of $N_{\hat{\theta}}$. Since $\mathcal{L}(V_\sigma) = M_{n_\sigma}(\mathbb{C}) = \mathcal{U}(n_\sigma)_\mathbb{C}$, then it follows that $\{T \in \mathcal{L}^{sk}(V_\sigma) | \text{Tr}(T) = 0\} = \mathcal{SU}(n_\sigma)_\mathbb{C}$ where $n_\sigma = \dim_\mathbb{C} V_\sigma$ and $\mathcal{U}(n_\sigma)_\mathbb{C}$, $\mathcal{SU}(n_\sigma)_\mathbb{C}$ are the complexifications of the Lie algebras of $\text{U}(n_\sigma)$, $\text{SU}(n_\sigma)$ respectively. We then deduce that

$$\mathcal{U}(n_\sigma)_\mathbb{C} = \mathbb{C}I_{n_\sigma} \oplus \mathcal{SU}(n_\sigma)_\mathbb{C}$$

Therefore it implies that

$$\hat{\Xi}_\sigma = M_\theta \oplus_{M_\theta} (N_{\hat{\theta}} \otimes \mathcal{SU}(n_\sigma)_\mathbb{C})^{\hat{\theta} \otimes \text{Ad}(\sigma)(G)}$$

Let us define

$$\Gamma(\text{ad}_\sigma(N_{\hat{\theta}})) = (N_{\hat{\theta}} \otimes \mathcal{U}(n_\sigma)_\mathbb{C})^{\hat{\theta} \otimes \text{Ad}(\sigma)(G)}.$$

Moreover, we put

$$\Omega^0(\text{ad}_\sigma(N_{\hat{\theta}})) = \Gamma(\text{ad}_\sigma(N_{\hat{\theta}})), \quad \Omega^1(\text{ad}_\sigma(N_{\hat{\theta}})) = \Omega^1(M_\theta, \Gamma(\text{ad}_\sigma(N_{\hat{\theta}})))$$

, and

$$\Omega_-^2(\text{ad}_\sigma(N_{\hat{\theta}})) = P_- \Omega^2(M_\theta, \Gamma(\text{ad}_\sigma(N_{\hat{\theta}}))).$$

We then introduce an inner product $\langle \cdot | \cdot \rangle$ on $\Omega^j(\text{ad}_\sigma(N_{\hat{\theta}}))$ ($j = 0, 1, 2$) by

$$\langle \omega | \eta \rangle = \int_{M_\theta} \text{tr}(*_\theta(\omega^* *_\theta \eta))$$

where $*_\theta$ is the Hodge operation on $\Omega^*(M_\theta)$, and tr is the canonical trace on $M_{n_\sigma}(\mathbb{C})$. Using this inner product, we also induce the metric topology on $\mathcal{C}_\pm(\Xi_\sigma)$ by the following lemma:

Lemma 3.1(cf:[T]) Let $\nabla_\sigma \in \mathcal{C}_+(\Xi_\sigma)$. Then it follows that

$$\mathcal{C}_+(\Xi_\sigma) = \nabla_\sigma + \Omega_+^1(\text{ad}_\sigma(N_{\hat{\theta}}))$$

where $\Omega_+^1(\text{ad}_\sigma(N_{\hat{\theta}}))$ is the set of all $\omega \in \Omega^1(\text{ad}_\sigma(N_{\hat{\theta}}))$ satisfying the equation:

$$\hat{\nabla}^-(\omega) + (\omega^2)_- = 0,$$

and $(\omega^2)_- = P_-(\omega^2)$.

Proof. Let $\nabla \in \mathcal{C}_+(\Xi_\sigma)$ and put $\omega = \nabla - \nabla_\sigma$, then it implies by definition that $\omega \in \hat{\Xi}_\sigma$. Since $\nabla, \nabla_\sigma \in \mathcal{C}_+(\Xi_\sigma)$, it follows that

$$P_-[\nabla, [\nabla, X]] = P_-[F_\nabla, X] = [F_\nabla^-, X] = 0$$

for all $X \in \hat{\Xi}_\sigma$ respectively, and the similar statement holds for ∇_σ . We compute that

$$F_\nabla = F_{\nabla_\sigma} + [\nabla_\sigma, \omega] + \omega^2$$

, which implies the conclusion. Q.E.D.

As M is compact oriented, it has a spin^c structure. As the similar case as a spin structure, it follows from [CD·V] that there exists a noncommutative spin^c structure $S(c)_\theta$ of M_θ as a right M_θ -module induced by the given spin^c one $S(c)$ of M . Let $S(c)_\theta^\pm$ be the half spin structures of M_θ induced by the given half spin^c one $S(c)^\pm$ of M respectively. We use the same notations as their smooth sections on M_θ . Then the following statement is easily seen by their definition:

Lemma 3.2(cf:[T])

$$(1) \quad \Omega^1(M_\theta) = S(c)_\theta^\pm \otimes S(c)_\theta^\mp$$

$$(2) \quad M_\theta \oplus \Omega_\pm^2(M_\theta) = S(c)_\theta^\pm \otimes S(c)_\theta^\pm$$

respectively.

Proof. (1): By definition, $S(c)_\theta^\pm$ is the algebra of all even (odd) polynomials of $\Omega^1(M_\theta)$ with the Clifford multiplication c_θ respectively. Then $c_\theta(\Omega^1(M_\theta)S(c)_\theta^\pm) = S(c)_\theta^\mp$ respectively, which implies the conclusion. (2): The result is true if $\theta = 0$. This isomorphism can also be chosen $\tilde{T}^2$ -equivariantly, where $\tilde{T}^2$ is a covering group of T^2 . Then the statement follows. Q.E.D.

By Lemmas 3.1, $\mathcal{C}_+(\Xi_\sigma)$ is identified with $\Omega_+^1(ad_\sigma(N_{\hat{\theta}}))$ as an affine space respectively. We then show the next three lemma which seems to be quite useful showing our main result:

Lemma 3.3(cf:[T]) Let $T_\nabla(\mathcal{C}_+(\Xi_\sigma))$ be the tangent space of $\mathcal{C}_+(\Xi_\sigma)$ at $\nabla \in \mathcal{C}_+(\Xi_\sigma)$. Then it follows that

$$T_\nabla(\mathcal{C}_+(\Xi_\sigma)) = \{\omega \in \Omega^1(ad_\sigma(N_{\hat{\theta}})) \mid P_-[\nabla, \omega] = 0\}.$$

Proof. Let $\omega_t = \nabla_t - \nabla$ for a smooth curve $\nabla_t \in \mathcal{C}_+(\Xi_\sigma)$ with $\nabla_0 = \nabla$. Then it follows from definition that $\omega_t \in \Omega^1(ad_\sigma(N_{\hat{\theta}}))$.

We put ω the derivative of ω_t at $t = 0$ in $\Omega^1(ad_\sigma(N_{\hat{\theta}}))$. Then we see that

$$F_{\nabla_t} = F_{\nabla} + [\nabla, \omega_t] + \omega_t^2,$$

which deduces $F_{\nabla_t}'(0) = [\nabla, \omega]$ taking their derivatives at $t = 0$ since $\omega_0 = 0$. As $\nabla_t \in \mathcal{C}_+(\Xi_\sigma)$, it implies that $F_{\nabla_t} \in \Omega_+^2(ad(N_{\hat{\theta}}))$, which means that $P_-[\nabla, \omega] = 0$. Q.E.D.

Let $\Gamma(\Xi_\sigma)$ be the gauge group acting on the set $\mathcal{C}(\Xi_\sigma)$ of all compatible connections of Ξ_σ by $\gamma_u(\nabla) = u\nabla u^*$, ($u \in \Gamma(\Xi_\sigma, \nabla \in \mathcal{C}(\Xi_\sigma))$), and $\Gamma(\Xi_\sigma) \cdot \nabla$ the orbit of ∇ under $\Gamma(\Xi_\sigma)$. Then we show the following lemma:

Lemma 3.4(cf:[T]) Let $T_\nabla(\Gamma(\Xi_\sigma) \cdot \nabla)$ be the tangent space of $\Gamma(\Xi_\sigma) \cdot \nabla$ at $\nabla \in \mathcal{C}(\Xi_\sigma)$. Then we have that

$$T_\nabla(\Gamma(\Xi_\sigma) \cdot \nabla) = \{ [\nabla, X] \mid X \in \Omega^0(ad(N_{\hat{\theta}})) \}$$

for any $\nabla \in \mathcal{C}_+(\Xi_\sigma)$.

Proof. Let $X \in \Omega^0(ad_\sigma(N_{\hat{\theta}}))$ and put $\varphi_t = e^{-tX} \in \Gamma(\Xi_\sigma)$ for all $t \in \mathbb{R}$. Then we know that

$$\gamma_{\varphi_t}(\nabla)(\xi) = e^{-tX} \cdot \nabla(e^{tX}\xi) = t[\nabla, X](\xi) + \nabla(\xi)$$

for all $\xi \in \Xi_\sigma$. Taking their derivatives at $t = 0$, we have that

$$\left. \frac{d}{dt} \gamma_{\varphi_t}(\nabla) \right|_{t=0}(\xi) = [\nabla, X](\xi)$$

for all $\xi \in \Xi_\sigma$. Therefore the conclusion follows. Q.E.D.

As we know that $\Gamma(\Xi_\sigma) \cdot \nabla \subseteq \mathcal{C}_+(\Xi_\sigma)$ for any $\nabla \in \mathcal{C}_+(\Xi_\sigma)$, we may imagine by Lemma 3.3 and 3.4 the next corollary:

Corollary 3.5(cf:[T]) Let $\mathcal{M}_\pm(\Xi_\sigma)$ be the moduli space of $\mathcal{C}_\pm(\Xi_\sigma)$ by $\Gamma(\Xi_\sigma)$. Then the tangent space $T_{[\nabla]}(\mathcal{M}_+(\Xi_\sigma))$ of $\mathcal{M}_+(\Xi_\sigma)$ at $[\nabla] \in \mathcal{M}_+(\Xi_\sigma)$ is isomorphic to $T_\nabla(\mathcal{C}_+(\Xi_\sigma))/T_\nabla(\Gamma(\Xi_\sigma) \cdot \nabla)$ as a $\mathbb{C}$ -linear space.

Let us define $\hat{\nabla}^-(\omega) = P_-[\nabla, \omega]$ for all $\omega \in \Omega^1(ad_\sigma(N_{\hat{\theta}}))$. We then can show the following lemma which is well known in undeformed cases by [AHS] and in a special deformed case by [LS]:

Lemma 3.6(cf:[T])

$$0 \rightarrow \Omega^0(ad_\sigma(N_{\hat{\theta}})) \xrightarrow{\hat{\nabla}} \Omega^1(ad_\sigma(N_{\hat{\theta}})) \xrightarrow{\hat{\nabla}^-} \Omega_-^2(ad_\sigma(N_{\hat{\theta}})) \rightarrow 0$$

for $\nabla \in \mathcal{C}_+(\Xi_\sigma)$, where $\hat{\nabla}(X) = [\nabla, X]$ for $X \in \Omega^0(ad_\sigma(N_{\hat{\theta}}))$ and $\hat{\nabla}^-(\omega) = P_-[\nabla, \omega]$ for $\omega \in \Omega^1(ad_\sigma(N_{\hat{\theta}}))$.

Proof. We first show that $\hat{\nabla}^- \cdot \hat{\nabla} = 0$. Indeed, as $\nabla \in \mathcal{C}_+(\Xi_\sigma)$,

$$\hat{\nabla}^- \cdot \hat{\nabla}(X) = P_-[\nabla, [\nabla, X]] = P_-[F_\nabla, X] = [F_{\nabla^-}, X] = 0$$

for any $X \in \Omega^0(ad_\sigma(N_{\hat{\theta}}))$. We next show that $\text{Ker } \hat{\nabla} = 0$. In fact, suppose $X \in \text{Ker } \hat{\nabla}$. Since $\hat{\nabla}$ commutes with the action α of T^2 and $ad(N_{\hat{\theta}})$ has $\mathcal{U}(n_\sigma)_\mathbb{C}$ as fibres, then it follows that

$$\hat{\nabla} = d_\theta^n + \omega_n$$

for some $\omega_n \in M_n(\Omega^1(N_{\hat{\theta}}))$, where $(\omega_n)_{j,k}$ are all central in $\Omega^1(N_{\hat{\theta}})$, which means that $L_\theta(\omega_n) = \omega_n$. Hence it implies that $\hat{\nabla}(X) = P_\sigma \cdot d_\theta^n(X) = 0$. Since $X = L_\theta(X^0)$ for a $X^0 \in \Omega^0(ad(N))$, then we have that

$$\hat{\nabla}(X) = P_\sigma \cdot d_\theta^n(X) = L_\theta \cdot p_\sigma \cdot d_M(X^0) = 0,$$

where p_σ is the projection in $M_n(C^\infty(M))$ corresponding to the right $C^\infty(M)$ -module $\Xi = (C^\infty(N) \otimes V_\sigma)^{\otimes \sigma(G)}$ and d_M is the outer derivative of M . Then $p_\sigma \cdot d_M(X^0) = 0$, which implies that $X^0 = 0 \in \Omega^0(ad(N))$ by the irreducibility of $p_\sigma \cdot d_M$ on Ξ . This means that $\text{Ker } \hat{\nabla} = 0$. Q.E.D.

Let us define $\mathbb{H}_-^1 = \text{Ker } \hat{\nabla}^- / \text{Im } \hat{\nabla}$ and $\mathbb{H}_-^2 = \Omega_-^2(ad_\sigma(N_{\hat{\theta}})) / \text{Im } \hat{\nabla}^-$. It follows from Lemma 3.5 that $\mathbb{H}_-^1$ is isomorphic to $T_{[\nabla]}(\mathcal{M}_+(\Xi_\sigma))$ as a $\mathbb{C}$ -linear space for any $\nabla \in \mathcal{C}_+(\Xi_\sigma)$. Moreover, we introduce the Laplace type M_θ -operators Δ_j^- on $\Omega^j(ad_\sigma(N_{\hat{\theta}}))$ for $j = 1, 2$ as follows: let $\Delta_j^- = (\hat{\nabla}^-)^* \cdot \hat{\nabla}^- + \hat{\nabla}^- \cdot (\hat{\nabla}^-)^*$ on $\Omega^1(ad_\sigma(N_{\hat{\theta}}))$ and $\Delta_j^- = \hat{\nabla}^- \cdot (\hat{\nabla}^-)^*$ on $\Omega^2(ad_\sigma(N_{\hat{\theta}}))$. Then we easily observe the following lemma:

Lemma 3.7(cf:[T]) $\mathbb{H}_-^j$ are isomorphic to $\text{Ker } \Delta_j^-$ ($j = 1, 2$) as a right M_θ -module respectively, and they are finitely generated projective right M_θ -modules.

Proof. $\mathbb{H}_-^1$ is M_θ -isomorphic to $\text{Ker } \hat{\nabla}^- \cap \text{Ker } (\hat{\nabla}^-)^*$, which is equal to $\text{Ker } \Delta_j^-$. The similar way is also valid for $\mathbb{H}_-^2$. As Δ_j^- are elliptic, the rest is well known. Q.E.D.

By Lemma 3.6, the elliptic complex can be described by the following generalized signature M_θ -operator:

$$\hat{\nabla}^- + (\hat{\nabla}^-)^* : \Omega^1(ad_\sigma(N_{\hat{\theta}})) \rightarrow \Omega^0(ad_\sigma(N_{\hat{\theta}})) \oplus \Omega_-^2(ad_\sigma(N_{\hat{\theta}}))$$

for all $\nabla \in \mathcal{C}_+(\Xi_\sigma)$. By Lemma 3.7, $\mathbb{H}_-^j$ induce the $\mathbb{K}_0(M_\theta)$ -element $[\mathbb{H}_-^j]$ ($j = 1, 2$) whose K-theoretic ranks are finite. Let us define the K-theoretic index of $\hat{\nabla}^- + (\hat{\nabla})^*$ as follows:

$$\text{Index}_{M_\theta}(\hat{\nabla}^- + (\hat{\nabla})^*) = [\text{Ker}(\hat{\nabla}^- + (\hat{\nabla})^*)] - [\text{Coker}(\hat{\nabla}^- + (\hat{\nabla})^*)]$$

Using Lemma 3.6, we then obtain the following lemma which is a noncommutative version of Atiyah-Singer index theorem due to Connes [C] (cf:[MS]):

Lemma 3.8(cf:[T])

$$\text{Index}_{M_\theta}(\hat{\nabla}^- + (\hat{\nabla})^*) = [\mathbb{H}_-^1] - [\mathbb{H}_-^2]$$

Let $[M_\theta]$ be the fundamental cyclic cocycle of M_θ which is essentially defined by the JLO-cocycle appeared in [C]. Then it follows by Lemma 3.8 that

Corollary 3.9

$$\langle [M_\theta], \text{Index}_{M_\theta}(\hat{\nabla}^- + (\hat{\nabla})^*) \rangle = \text{rank}_{M_\theta}[\mathbb{H}_-^1] - \text{rank}_{M_\theta}[\mathbb{H}_-^2]$$

On the other hand, we consider the Dirac type M_θ -operator D_∇^+ on $\Omega^1(ad_\sigma(N_{\hat{\theta}}))$ associated with $\nabla \in \mathcal{C}_+(\Xi_\sigma)$ defined by the following process: By Lemma 3.2.(1),

$$\begin{aligned} \Omega^1(ad_\sigma(N_{\hat{\theta}})) &= \Omega^0(ad_\sigma(N_{\hat{\theta}})) \otimes (S(c)_\theta^+ \otimes S(c)_\theta^-) \\ &\xrightarrow{\hat{\nabla}} \Omega^0(ad_\sigma(N_{\hat{\theta}})) \otimes \Omega^1(M_\theta) \otimes (S(c)_\theta^+ \otimes S(c)_\theta^-) \\ &\xrightarrow{\text{Id} \otimes c_\theta^+} \Omega^0(ad_\sigma(N_{\hat{\theta}})) \otimes (S(c)_\theta^- \otimes S(c)_\theta^-) = \Omega^0(ad_\sigma(N_{\hat{\theta}})) \oplus \Omega_-^2(ad_\sigma(N_{\hat{\theta}})) \end{aligned}$$

Lemma 3.10(cf:[T]) Given a $\nabla \in \mathcal{C}_+(\Xi_\sigma)$,

$$\text{Index}_{M_\theta}(D_\nabla^+) = \text{Index}_{M_\theta}(\hat{\nabla}^* + \hat{\nabla}^-)$$

in $K_0(M_\theta)$.

Proof. Since $\Omega(M_\theta) \simeq \Omega(M)$ as vector spaces and $\hat{\nabla}$ and $*$ -operation commute with T^2 -action α , then the operator $\hat{\nabla}^* + \hat{\nabla}^-$ can be shifted to the elliptic operator $(d_M)^* + p_- \cdot d_M$ from $\Omega^1(M)$ to $\Omega^0(M) \oplus \Omega_-^2(M)$. Let P_-^0 be the projection from $\Omega^2(M)$ to $\Omega_-^2(M)$ respectively. Using the same argument as in [AHS] and taking their

principal symbols, it follows from Lemma 3.2 that $(d_M)^* + P_{\mp}^0 \cdot d_M$ is identified with the Dirac operator D^+ in the following:

$$D^+ : S(c)^+ \otimes S(c)^- \rightarrow S(c)^- \otimes S(c)^-$$

, so far taking their $C^\infty(M)$ -indices. Then $\hat{\nabla}^* + \hat{\nabla}^-$ is identified with D_σ^+ defined before:

$$D_\sigma^+ : \Omega^0(ad_\sigma(N_{\hat{\theta}})) \otimes (S(c)_\theta^+ \otimes S(c)_\theta^-) \rightarrow \Omega^0(ad_\sigma(N_{\hat{\theta}})) \otimes (S(c)_\theta^- \otimes S(c)_\theta^-)$$

so far taking their M_θ -indices. This implies the conclusion. Q.E.D.

In what follows, we want to determine the geometric structure of the moduli space $M_+(\Xi_\sigma)$ of $C_+(\Xi_\sigma)$ by the gauge group $\Gamma(\Xi_\sigma)$. First of all, we introduce a noncommutative Kuranishi map in the following process : Given a $\nabla \in \mathcal{C}_+(\Xi_\sigma)$, we define a Sobolev norm $\|\cdot\|_p$, ($p \gg 4$) on $\Omega_+^1(ad_\sigma(N_{\hat{\theta}}))$ by

$$\|\omega\|_p = \sum_{k=0}^p \int_{M_\theta} \|(\hat{\nabla})^k(\omega)\|^2$$

for $\omega \in \Omega_+^1(ad_\sigma(N_{\hat{\theta}}))$. Let S_+^p be the completion of $\Omega_+^1(ad_\sigma(N_{\hat{\theta}}))$ with respect to $\|\cdot\|_p$. By their definition, $\hat{\nabla}$, $\hat{\nabla}^*$ are bounded from S_+^p to S_+^{p+1} , S_+^{p-1} respectively.

Since Δ_j^- is elliptic in $\text{End}_{M_\theta}(S_+^p)$, it is a M_θ -Fredholm operator on S_+^p as well. Let $P_{\mathbb{H}_-^2}$ be the projection on $\mathbb{H}_-^2$, and define the Green M_θ -operator G^- of Δ_2^- on S_+^p as follows :

$$G^- = (\text{Id}_{S_+^p} - P_{\mathbb{H}_-^2}) \cdot (\Delta_2^-)^{-1} = (\Delta_2^-)^{-1} \cdot (\text{Id}_{S_+^p} - P_{\mathbb{H}_-^2}).$$

Then we easily see that

- (1) : $\text{Im } G^- = \text{Im } \Delta_2^-$,
- (2) : $G^- \cdot \Delta_2^- = \Delta_2^- \cdot G^-$,
- (3) : $G^- \cdot \Delta_2^- = \text{Id}_{S_+^p} - P_{\mathbb{H}_-^2}$

We now introduce a densely defined map Π_{∇}^- on S_+^p by

$$\Pi_{\nabla}^-(\omega) = \omega + (\hat{\nabla}^-)^* \cdot G^\pm \{(\omega^2)_-\}$$

for all $\omega \in \Omega_+^1(ad_\sigma(N_{\hat{\theta}}))$ respectively, which is called a noncommutative Kuranishi map, which is known in commutative cases

(cf:[AHS]). Then we deduce the following observation:

Lemma 3.11(cf:[T])

$$\Omega_+^1(ad_\sigma(N_{\hat{\theta}})) = \text{Ker } (\hat{\nabla}^- \cdot \Pi_{\nabla}^-) \cap \{P_{\mathbb{H}_-^2}((\omega^2)_-) = 0\} \ .$$

Proof. By Lemma 3.1, $\omega \in \Omega_+^1(ad_\sigma(N_{\hat{\theta}}))$ satisfies

$$\hat{\nabla}^-(\omega) + (\omega^2)_- = 0 \ .$$

Then we compute

$$\begin{aligned} \hat{\nabla}^- \cdot \Pi_{\nabla}^-(\omega) &= \hat{\nabla}^-(\omega) + \Delta_2^- \cdot G^-((\omega^2)_-) \\ &= \hat{\nabla}^-(\omega) + (\omega^2)_- - P_{\mathbb{H}_-^2}((\omega^2)_-) = -P_{\mathbb{H}_-^2}((\omega^2)_-) \end{aligned}$$

, which implies that

$$\hat{\nabla}^- \cdot \Pi_{\nabla}^-(\omega) = -P_{\mathbb{H}_-^2}((\omega^2)_-) \ .$$

By the definition of $\mathbb{H}_-^2$, it follows that

$$(\hat{\nabla}^-)^* \cdot \hat{\nabla}^- \cdot \Pi_{\nabla}^-(\omega) = 0$$

, which implies by taking their inner products that

$$\hat{\nabla}^- \cdot \Pi_{\nabla}^-(\omega) = P_{\mathbb{H}_-^2}((\omega^2)_-) = 0 \ .$$

The converse implication is also easily seen. Q.E.D.

We then observe together with Lemma 3.1 the following corollary:

Corollary 3.12 If $\nabla \in \mathcal{C}_+(\Xi_\sigma)$, then it follows that

$$\mathcal{C}_+(\Xi_\sigma) = \nabla + \text{Ker } (\hat{\nabla}^- \cdot \Pi_{\nabla}^-) \cap \{P_{\mathbb{H}_-^2}((\omega^2)_-) = 0\} \ .$$

It also follows from Lemma 3.11 that

Corollary 3.13 Suppose $\mathbb{H}_\mp^2 = 0$ for a $\nabla \in \mathcal{C}_+(\Xi_\sigma)$, then it follows that

$$\Omega_+^1(ad_\sigma(N_{\hat{\theta}})) = \text{Ker } (\hat{\nabla}^- \cdot \Pi_{\nabla}^-)$$

Moreover, we obtain by using Lemma 3.11 the following lemma:

Lemma 3.14(cf:[T])

$$\Pi_{\nabla}^-(\Omega_+^1(ad_\sigma(N_{\hat{\theta}}))) = \mathbb{H}_-^1$$

for all $\nabla \in \mathcal{C}_+(\Xi_\sigma)$,

Proof. As $\hat{\nabla}^- \cdot \hat{\nabla} = 0$ for all $\nabla \in \mathcal{C}_+(\Xi_\sigma)$, then $(\hat{\nabla})^* \cdot (\hat{\nabla}^-)^* = 0$. Let $\eta = \Pi_{\nabla}^-(\omega)$ for any $\omega \in \Omega_+^1(ad_\sigma(N_{\hat{\theta}}))$. Then we have that

$$(\hat{\nabla})^*(\eta) = (\hat{\nabla})^*(\omega) + \Delta_2^- \cdot G^-(\omega^2)_- = 0$$

, respectively. By Lemma 3.11,

$$\hat{\nabla}^-(\eta) = P_{\mathbb{H}_-^2}((\omega^2)_-) = 0.$$

Since $\text{Ker } (\hat{\nabla})^* = \Omega^1(ad_\sigma(N_{\hat{\theta}}))$, then the conclusion follows. Q.E.D.

Using the Sobolev space S_+^p , we compute the Frechet differentiation $\left. \frac{d\Pi_{\nabla}^-}{dt} \right|_{t=0}$ of Π_{∇}^- on $\Omega_+^1(ad_\sigma(N_{\hat{\theta}}))$ in the following lemma:

Lemma 3.15

$$\left. \frac{d\Pi_{\nabla}^-}{dt} \right|_{t=0} = \text{Id}$$

on $\Omega_+^1(ad_\sigma(N_{\hat{\theta}}))$ with respect to $\|\cdot\|_p$.

Proof.

$$\begin{aligned} \left. \frac{d\Pi_{\nabla}^-}{dt} \right|_{t=0}(\omega) &= \lim_{t \rightarrow 0} t^{-1}(\Pi_{\nabla}^-(t\omega) - \Pi_{\nabla}^-(0)) \\ &= \lim_{t \rightarrow 0} (\omega + t(\hat{\nabla}^-)^* \cdot G^-(\omega^2)_-) = \omega \end{aligned}$$

for any $\omega \in \Omega_+^1(ad_\sigma(N_{\hat{\theta}}))$, which implies the conclusion. Q.E.D.

By the inverse function theorem on Banach spaces, there exists a $\epsilon > 0$ neighborhood $U(0, \epsilon)$ of $0 \in S_+^p$ with respect to $\|\cdot\|_p$ on which $(\Pi_{\nabla}^-)^{-1}$ is diffeomorphic. Using this fact, we next show the following lemma:

Lemma 3.16 $U(0, \epsilon) \cap \mathbb{H}_-^1$ is diffeomorphic to $(\Pi_{\nabla}^-)^{-1}(U(0, \epsilon)) \cap \Omega_+^1(ad_\sigma(N_{\hat{\theta}}))$ under $(\Pi_{\nabla}^-)^{-1}$.

Proof. By Lemma 3.14, we know that

$$\Pi_{\nabla}^-(\Omega_+^1(ad_\sigma(N_{\hat{\theta}}))) = \mathbb{H}_-^1,$$

which implies the conclusion. Q.E.D.

Let $P\Gamma(\Xi_\sigma) = \Gamma(\Xi_\sigma)/Z(\Gamma(\Xi_\sigma))$ be the projective gauge group of $\Gamma(\Xi_\sigma)$ where $Z(\Gamma(\Xi_\sigma))$ is the center of $\Gamma(\Xi_\sigma)$. Since Ξ_σ is an irreducible right M_θ -module, $Z(\Gamma(\Xi_\sigma))$ is actually the center $ZU(M_\theta)$ of the unitary group $U(M_\theta)$ of M_θ . We define the gauge action γ of $P\Gamma(\Xi_\sigma)$ on $\mathcal{C}_+(\Xi_\sigma)$ by $\gamma_{[u]}(\nabla)(\xi) = u\nabla u^*(\xi)$ for any $[u] = uZU(M_\theta) \in P\Gamma(\Xi_\sigma)$, $\nabla \in \mathcal{C}_+(\Xi_\sigma)$. We then easily see that:

Lemma 3.17 The action γ is effective.

Proof. The statement is almost clear by its definition. Q.E.D.

As the similar way to the commutative cases (cf:[AHS],[FU]), we then show the so-called slice theorem in noncommutative setting:

Lemma 3.18(cf:[T]) Let ϵ be in Lemma 3.16. Then there exist $0 < \delta \leq \epsilon$, $U(0, \delta) = \{\omega \in \Omega^1(ad_\sigma(N_{\hat{\theta}})) \mid \|\omega\|_p < \delta\}$, and a mapping

$$\psi : \nabla + U(0, \delta) \rightarrow P\Gamma(\Xi_\sigma)$$

with the property that

$$(1) : \hat{\nabla}^*(\gamma_{\psi(\nabla+\omega)}^*(\nabla + \omega) - \nabla) = 0$$

(2): there exists a $P\Gamma(\Xi_\sigma)$ -equivariant diffeomorphism

$$\Phi : \nabla + U(0, \delta) \rightarrow U(0, \text{Id}) \subseteq \text{Ker } \hat{\nabla}^* \times P\Gamma(\Xi_\sigma).$$

defined by

$$\Phi(\nabla + \omega) = (\gamma_{\psi(\nabla+\omega)}^*(\nabla + \omega) - \nabla, \psi(\nabla + \omega))$$

Proof. Let us consider the next equation defined by

$$L_{\hat{\nabla}}(\omega, [u]) = \hat{\nabla}^*(u^*\nabla u + u^*\omega u) = 0$$

for $\nabla + \omega \in \mathcal{C}_+(\Xi_\sigma)$ and $[u] \in P\Gamma(\Xi_\sigma)$. Then the differential $\delta L_{\hat{\nabla}}$ of $L_{\hat{\nabla}}$ at $(0, \text{Id})$ is given by

$$\delta L_{\hat{\nabla}} : \Omega_+^1(ad_\sigma(N_{\hat{\theta}})) \times \Omega^0(ad_\sigma(N_{\hat{\theta}})) \rightarrow \Omega^0(ad_\sigma(N_{\hat{\theta}}))$$

$$\delta\omega \oplus \delta[u] \rightarrow \hat{\nabla}^*([\nabla, \delta u] + \delta\omega)$$

, The partial differential $\delta_2 L_{\hat{\nabla}}$ of $\delta L_{\hat{\nabla}}$ in the second factor is $\hat{\nabla}^* \cdot \hat{\nabla}$, which is a self-adjoint elliptic operator. Since $\text{Ker } \hat{\nabla} = 0$, then it follows by standard elliptic theory $\hat{\nabla}^* \cdot \hat{\nabla}$ is invertible on S_+^p . By the implicit function theorem in Hilbert space, we obtain a neighbourhood

$U(\omega, \delta)$ with $0 < \delta \leq \epsilon$ and a map $\psi : U(\omega, \delta) \rightarrow P\Gamma(\Xi_\sigma)$ satisfying our conditions. Hence we obtain the diffeomorphism Φ on $\nabla + U(0, \delta)$ cited in this lemma. Since we see that

$$L_{\hat{\nabla}}([v]^*\omega, [v^{-1}u]) = L_{\hat{\nabla}}(\omega, [u]) ,$$

for all $[u], [v] \in P\Gamma(\Xi_\sigma)$, $\omega \in \Omega^1(ad_\sigma(N_{\hat{\theta}}))$, then Φ is chosen $P\Gamma(\Xi_\sigma)$ -equivariantly. Q.E.D.

By (2) of the above lemma 3.18, $U(0, \delta)$ can be chosen as $P\Gamma(\Xi_\sigma)$ -invariant. We then prove the following lemma:

Lemma 3.19(cf:[T]) Let $\nabla \in C_+(\Xi_\sigma)$. Then there exists a $\delta > 0$ such that

$$\nabla_\sigma + \{U(0, \delta) \cap \Omega_+^1(ad_\sigma(N_{\hat{\theta}})) \cap \text{Ker } (\hat{\nabla})^*\}$$

is diffeomorphically imbedded in the moduli space $\mathcal{M}_+(\Xi_\sigma)$ of $\mathcal{C}_+(\Xi_\sigma)$ by $\Gamma(\Xi_\sigma)$.

Proof. By Lemma 3.18, $\nabla_\sigma + \{U(0, \delta) \cap \Omega_+^1(ad_\sigma(N_{\hat{\theta}}))\}$ is $P\Gamma(\Xi_\sigma)$ -equivariantly diffeomorphic to $U(0, \text{Id}) \subseteq \text{Ker } \hat{\nabla}^* \times P\Gamma(\Xi_\sigma)$. Then we conclude that

$$(\nabla_\sigma + \{U(0, \delta) \cap \text{Ker } (\hat{\nabla})^*\}) \cap C_+(\Xi_\sigma)$$

is diffeomorphically embedded in $\mathcal{M}_+(\Xi_\sigma)$. Q.E.D.

We next show the following lemma (cf:[FU]):

Lemma 3.20 $\mathcal{M}_+(\Xi_\sigma)$ is a Hausdorff space.

Proof. It suffices to show that $\mathcal{M}(\Xi_\sigma) = \mathcal{C}(\Xi_\sigma)/\Gamma(\Xi_\sigma)$ is Hausdorff. So we need to check that

$$\Upsilon = \{(\nabla, \gamma_u(\nabla)) \mid \nabla \in \mathcal{C}(\Xi_\sigma), u \in \Gamma(\Xi_\sigma)\}$$

is closed in $\mathcal{C}(\Xi_\sigma) \times \mathcal{C}(\Xi_\sigma)$. In fact, if $\{(\nabla + \omega_j, \gamma_{u_j}(\nabla + \omega_j))\}_j \subseteq \Upsilon$ is convergent to $(\nabla + \omega, \nabla + \omega')$, then $\omega_j \rightarrow \omega$ and

$$\gamma_{u_j}(\nabla + \omega_j) = \nabla + u_j \hat{\nabla}(u_j^*) + u_j \omega u_j^* \rightarrow \nabla + \omega' .$$

Put $\omega'_j = u_j \hat{\nabla}(u_j^*) + u_j \omega u_j^* \in \Omega^1(ad_\sigma(N_{\hat{\theta}}))$. Then $\hat{\nabla}(u_j^*) = u_j^* \omega' - \omega u_j^*$. Since $\|u_j \nabla\|_k$ ($0 \leq k \leq p-1$) are bounded by definition of the Sobolev norms $\|\cdot\|$, Hence $\{u_j\}$ is bounded in $S^p(\Omega^0(ad_\sigma(N_{\hat{\theta}})))$. Then it follows by Rellich's theorem that there exists a subsequence

$\{u_k\}$ of $\{u_j\}$ such that $\{u_k\}$ is convergent in $S^{p-1}(\Omega^1(ad_\sigma(N_{\hat{\theta}})))$. Hence $u_k^* \omega' - \omega u_k^*$ converges in $S^{p-1}(\Omega^1(ad_\sigma(N_{\hat{\theta}})))$, so is $\{\hat{\nabla}(u_k)\}_k$ in $S^{p-1}(\Omega^1(ad_\sigma(N_{\hat{\theta}})))$ as well. Therefore $\{u_k\}_k$ converges to $u \in S^p(\Omega^0(ad_\sigma(N_{\hat{\theta}})))$. Since $\Gamma(\Xi_\sigma)$ is closed in $S^p(\Omega^0(ad_\sigma(N_{\hat{\theta}})))$, we conclude that $\gamma_u(\nabla + \omega) = \nabla + \omega'$. This completes the proof. Q.E.D.

We now state our main theorem which is a generalization of the one appeared in [T] as follows:

Theorem 3.21 Let M be a compact oriented toric 4-dimensional Riemannian manifold, and $G \xrightarrow{\varrho} P \rightarrow M$ a principal G -bundle over M , where G is a compact connected Lie group. Suppose $G \xrightarrow{\tilde{\varrho}} P_{\tilde{\theta}} \rightarrow M_\theta$ exists, then given the finite generated projective irreducible right M_θ -module $\Xi_\sigma = (P_\theta \otimes V_\sigma)^{\tilde{\varrho} \otimes Ad(\sigma)(G)}$ for a highest weight σ of G , the moduli space $\mathcal{M}_+(\Xi_\sigma)$ of $\mathcal{C}_+(\Xi_\sigma)$ under $\Gamma(\Xi_\sigma)$ is a locally smooth manifold with its dimension:

$$< [M_\theta], \text{ch}_\theta(\Xi_\sigma \otimes S(c)^-) > + \text{rank}_{M_\theta} [\mathbb{H}_-^2]$$

where ch_θ is the Connes-Chern character from $K_0(M_\theta)$ to the periodic cyclic homology of M_θ , and $S(c)^-$ is the smooth sections of the $(-)$ -half spin^c -structure of M .

Proof. By Corollary 3.9, Lemma 3.17, 3.19 and 3.20, $\mathcal{M}_+(\Xi_\sigma)$ is a smooth manifold. By Lemma 3.16, its dimension is equal to $\text{rank}_{M_\theta} \mathbb{H}_-^1$, which is by Corollary 3.9 and Lemma 3.10 that

$$< [M_\theta], \text{ch}_\theta(\Xi_\sigma \otimes S(c)^-) > + \text{rank}_{M_\theta} [\mathbb{H}_-^2] .$$

This completes the proof. Q.E.D.

noindent From this theorem, we deduce several useful corollaries which are well known in the case of ordinary manifolds as well as a noncommutative 4-sphere case:

Corollary 3.22 In the above theorem, suppose $[\mathbb{H}_-^2] = 0 \in K_0(M_\theta)$ for all $\nabla \in \mathcal{C}_+(\Xi_\sigma)$, then the instanton moduli space $\mathcal{M}_+(\Xi_\sigma)$ of Xi_σ is a smooth manifold with its dimension:

$$< [M_\theta], \text{ch}_\theta(\Xi_\sigma \otimes S(c)^-) > .$$

Proof. By Corollary 3.12 and Theorem 3.21, the conclusion follows. Q.E.D.

Corollary 3.23 Let M be a compact oriented toric Riemannian 4-manifold with positive scalar curvature, and G a compact connected Lie group. Suppose there exists a noncommutative principal G -bundle

$$G \xrightarrow{\tilde{g}} N_{\tilde{g}} \rightarrow M_{\theta} ,$$

then given a highest weight σ of G and consider the right M_{θ} -module $\Xi_{\sigma} = (N_{\tilde{g}} \otimes V_{\sigma})^{\tilde{g} \otimes \text{Ad}(\sigma)(G)}$, then $\mathcal{M}_+(\Xi_{\theta})$ is a smooth manifold with dimension:

$$\dim \mathcal{M}_+(\Xi_{\sigma}) = \langle [M_{\theta}], ch_{\theta}(\Xi_{\theta} \otimes S(c)^{-} \rangle$$

Proof. By the same method as in [AHS], as the manifold M has a positive scalar curvature, we have that $(\mathbb{H}^0)_-^2 = 0$ for the 2-cohomology $(\mathbb{H}^0)_-^2$ with respect to the Laplacian associated with $(d_M)^* + P_-^0 \cdot d_M$. Since the quantization map L_{θ} from $C^{\infty}(M)$ to M_{θ} can be lifted to a K-theoretic isomorphism (cf:[R]) and $L_{\theta} \cdot d_M = d_{\theta} \cdot L_{\theta}$, then it implies that $\mathbb{H}_-^2 = 0$ for all $\nabla \in \mathcal{C}_+(\Xi_{\sigma})$. Hence the conclusion follows from Corollary 3.22. Q.E.D.

In what follows, we compute the typical example using the main theorem obtained in §3 :

§4. Example In what follows, we compute the typical example using the main theorem obtained in §3. Let us consider the Hopf bundle:

$$U(1) \longrightarrow S^5 \longrightarrow \mathbb{CP}^2 ,$$

and its associated noncommutative one:

$$U(1) \xrightarrow{\varrho} S_{\theta}^5 \longrightarrow \mathbb{CP}_{\theta}^2$$

Let 1 be the highest weight of $U(1)$ and $\Xi_1 = (S_{\theta}^5 \otimes \mathbb{C})^{\varrho \otimes \sigma_1^{-1}}(U(1))$ be its associated right $\mathbb{CP}_{\theta}^2$ -module where σ_1 is the character of $U(1)$ corresponding to 1. Let z_j ($j = 1, 2, 3$) be the 3-generators of S_{θ}^5 and put $P_1 = (z_1^*, z_2^*, z_3^*)^t (z_1, z_2, z_3)$ be the projection of $M_3(\mathbb{CP}_{\theta}^2)$ since $(z_1, z_2, z_3)(z_1^*, z_2^*, z_3^*)^t = I$. Then $\Xi_1 = P_1(\mathbb{CP}_{\theta}^2)^3$ up to isomorphism. Let $\nabla = P_1 \cdot d_{\theta}^3$ the Grassmann connection of Ξ_1 . Then $\nabla \in \mathcal{C}_+(\Xi_1)$. Now we compute the Connes-Chern character $ch_{\theta}(\Xi_1)$ of Ξ_1 . Then it follows from [C] (cf:[M]) that

$$ch_0(\Xi_1) = Tr(P_1) = \sum_j z_j^* z_j = I \in \mathbb{CP}_{\theta}^2 ,$$

$$ch_1(\Xi) = Tr((P - \frac{1}{2}I) \otimes P^{\otimes 2}) , \quad ch_2(\Xi_1) = Tr((P - \frac{1}{2}I) \otimes P^{\otimes 4})$$

where Tr is the canonical trace of $M_3(\mathbb{C})$. We also compute the chern character $ch(S(c)^-)$ of the spin^c bundle $S(c)^-$ over $\mathbb{CP}^2$. Let us take the real Fubini-Study metric as follows:

$$ds^2 = \frac{dr^2 + r^2 e_z^2}{(1+r^2)^2} + \frac{r^2(e_x^2 + e_y^2)}{(1+r^2)}$$

where $\rho_1 = r \cos(\psi) \exp(i\alpha)$, $\rho_2 = r \sin(\psi) \exp(i\beta)$ for any $(\rho_1, \rho_2, 1) \in \mathbb{CP}^2$,

and

$$\begin{aligned} e_x &= -\sin(\phi) \cos(\phi) \cos(\alpha + \beta) d\alpha + \sin(\phi) \cos(\phi) \cos(\alpha + \beta) d\beta \\ &\quad + \sin(\alpha + \beta) d\psi, \\ e_y &= -\sin(\phi) \cos(\phi) \cos(\alpha + \beta) d\alpha + \sin(\phi) \cos(\phi) \cos(\alpha + \beta) d\beta \\ &\quad - \cos(\alpha + \beta) d\psi, \\ e_z &= \cos^2(\phi) d\alpha + \sin^2(\psi) d\beta. \end{aligned}$$

We then define the spin^c connection ∇ on the smooth sections of $S(c)^-$ as follows:

$$\nabla = \left(-\frac{1}{4} \sum_{j,k=0,1,2,3} \omega_{jk} \Gamma^{jk} + iI\right)$$

where

$$\Gamma^0 = \begin{pmatrix} I_2 & 0 \\ 0 & -I_2 \end{pmatrix}, \quad \Gamma^j = \begin{pmatrix} 0 & i\sigma^j \\ i\sigma^j & 0 \end{pmatrix}, \quad \Gamma^{jk} = \frac{1}{2}[\Gamma^j, \Gamma^k].$$

for the Pauli matrices σ^j ($j = 1, 2, 3, 4$), and

$$\begin{aligned} \omega_{00} &= 0, \quad \omega_{01} = \frac{-1}{(1+r^2)^{1/2}} e_x, \quad \omega_{02} = \frac{-1}{(1+r^2)^{1/2}} e_y, \quad \omega_{03} = \frac{r^2-1}{r^2+1} e_z, \\ \omega_{10} &= \frac{-1}{(1+r^2)^{1/2}} e_x, \quad \omega_{11} = 0, \quad \omega_{12} = \frac{1+2r^2}{1+r^2} e_z, \quad \omega_{13} = \frac{-1}{(1+r^2)^{1/2}} e_y, \\ \omega_{20} &= \frac{-1}{(1+r^2)^{1/2}} e_y, \quad \omega_{21} = \frac{1+2r^2}{1+r^2} e_z, \quad \omega_{22} = 0, \quad \omega_{23} = \frac{1}{(1+r^2)^{1/2}} e_x, \\ \omega_{30} &= \frac{r^2-1}{r^2+1} e_z, \quad \omega_{31} = \frac{-1}{(1+r^2)^{1/2}} e_y, \quad \omega_{32} = \frac{1}{(1+r^2)^{1/2}} e_x, \quad \omega_{33} = 0. \end{aligned}$$

Then we compute the curvature F_∇ of ∇ as follows:

$$F_{\nabla} = \nabla^2 = \frac{1}{8}(\sum_{k=1,2,3} \omega_{0k}\omega_{0k+1}\Gamma^{0k}\Gamma^{0k+1}) + \omega_{0k}\omega_{(k+2)k}\Gamma^{0k}\Gamma^{(k+2)k} \\ + \sum_{j=0,1,2,3} \omega_{j(j+1)}\omega_{(j+1)(j+2)}\Gamma^{j(j+1)}\Gamma^{(j+1)(j+2)} + \omega_{01}\omega_{31}\Gamma^{03}\Gamma^{31})$$

(mod.3). Then it follows by their definitions that

$$ch_0(S(c)^-) = 4, \quad Tr(F_{\nabla}) = 0, \quad ch_1(S(c)^-) = 0$$

Moreover, we deduce that

$$Tr(F_{\nabla}^2) = 0, \quad ch_2(S(c)^-) = 0$$

Hence we have that $ch(S(c)^-) = 4$. Then we compute by definition (cf:[C],[G·B],[M]) that

$$< [CP_{\theta}^2], ch_{\theta}(\Xi_1 \otimes S(c)^-) > = 4\{1 - \frac{1}{2}\varphi^1(1, 1, 1) - 6\varphi^2(1, 1, 1, 1, 1)\} = 4$$

where φ^k is the JLO-cyclic k -cocycles ($k = 1, 2$) of $\mathbb{CP}_{\theta}^2$. We next compute $\mathbb{H}_{-}^2$. First of all, we know that

$$\Gamma(ad_{\sigma_1}(S_{\theta}^5)) = (S_{\theta}^5 \otimes \mathbb{C})^{\varrho \otimes Ad(\sigma_1)(U(1))} = \mathbb{CP}_{\theta}^2$$

We also check the following:

$$\Omega_{-}^2(ad_{\sigma_1}(S_{\theta}^5)) = \Omega_{-}^2(\mathbb{CP}_{\theta}^2) = (\Omega_{-}^2(\mathbb{CP}^2) \otimes T_{\theta}^2)^{\phi_{*} \otimes \alpha^{-1}(T^2)}$$

,and

$$\text{Im } \hat{\nabla}_{-} = (\text{Im } d^{-} \otimes T_{\theta}^2)^{\phi_{*} \otimes \alpha^{-1}(T^2)}.$$

We then check that

$$\mathbb{H}_{-}^2 = \Omega_{-}^2(ad_{\sigma_1}(S_{\theta}^5))/\text{Im } \hat{\nabla}^{-} = ((\Omega_{-}^2(\mathbb{CP}^2)/\text{Im } d^{-}) \otimes T_{\theta}^2)^{\phi_{*} \otimes \alpha^{-1}(T^2)},$$

which is equal to $(\mathbb{H}_{-}^2(\mathbb{CP}^2) \otimes T_{\theta}^2)^{\phi_{*} \otimes \alpha^{-1}(T^2)}$. Since $\mathbb{CP}^2$ has a positive scalar curvature, it follows by the Bochner-Weitzenböck formula that $\mathbb{H}_{-}^2(\mathbb{CP}^2) = 0$. Therefore, we conclude that $\mathbb{H}_{-}^2 = 0$. We then obtain the following result:

Theorem 4.1 Let us take the following Hopf bundle:

$$U(1) \longrightarrow S^5 \longrightarrow \mathbb{CP}^2,$$

and its associated noncommutative one:

$$U(1) \xrightarrow{\varrho} S_{\theta}^5 \longrightarrow \mathbb{CP}_{\theta}^2$$

Let 1 be the highest weight of $U(1)$ and $\Xi_1 = (S_{\theta}^5 \otimes \mathbb{C})^{\varrho \otimes \sigma_1^{-1}}(U(1))$ be its associated right $\mathbb{CP}_{\theta}^2$ -module where σ_1 is the character of $U(1)$

corresponding to 1. Then its instanton moduli space $\mathcal{M}_+(\Xi_1)$ is a smooth 4-manifold.

Remark For the highest weight n of $U(1)$, let Ξ_n be its corresponding right $\mathbb{CP}_\theta^2$ -module. Then the instanton moduli space $\mathcal{M}_+(\Xi_n)$ is a smooth $4n$ -manifold using the similar method as for Ξ_1 .

References

- [AHS] M.F.Atiyah, N.J.Hitchin and I.M.Singer, Self duality in 4-dimensional Riemannian geometry, Proc.R.S.London A362 (1978), 425-461.
- [C] A.Connes, Noncommutative Geometry, Academic Press (1994).
- [CD·V] Connes and M.Dubois·Violette, Noncommutative Finite-Dimensional Manifolds. Spherical Manifolds and Related Examples, Comm.Math.Phys., 230 (2002), 539-579.
- [CL] A.Connes and G.Landi, Noncommutative manifolds: The instanton algebra and isospectral deformations, Comm.Math. Phys., 221 (2001), 141-159.
- [FU] D.S.Freed and K.K.Uhlenbeck, Instanton and 4-Manifolds, MSRI.Publ.1, Springer-Verlag (1984).
- [G·BVF] J.M.Gracia·Bondia, J.C.Varilly and H.Figueroa, Elements of Noncommutative Geometry, Birkhauser Advanced Texts (2001).
- [LS] G.Landi and W.van Suijlekom, Noncommutative instantons from twisted conformal symmetries, arXiv.math QA/0601554 (2006).
- [MS] C.C.Moore and C.Schochet, Grobal Analysis on Foliated Spaces, MSRI.Publ.,9 Springer-Verlag (1988).
- [M] J.W.Morgan, The Seiberg-Witten equations and applications to topology of smooth four-manifolds, Princeton University Press (1996).

[T] H.Takai, Moduli Spaces of Instantons on Noncommutative 4-Manifolds, arXiv math.DG/0610536 (2006).